

## ORAL ABSTRACT

# Laboratory Analysis of HIV Infections in HPTN 083: Injectable CAB for PrEP

Marzinke M, Grinsztejn B, Fogel J, Piwowar-Manning EM, Hanscom B, Coelho L, Cohen MS, Rinehart AR, Rooney JF, Adeyeye A, Anderson P, McCauley M, Landovitz R, Eshleman S, for the HPTN 083 Study Team

Presenting Author: Raphael J. Landovitz  
*University of California San Francisco*




### Disclosure:

Raphael J. Landovitz has served on scientific advisory boards for Gilead Sciences and Merck Inc, and has received honoraria from Roche, Inc and Janssen.




# HPTN 083 Study Design

livestream.com befindet sich jetzt im Vollbildmodus. [Vollbild beenden \(Esc\)](#)





# Pre-specified HIV Testing




**HPTN Laboratory Center testing (retrospective)**

Visits with reactive/positive site tests



**Back-testing**



**Blinded adjudication of study endpoints**



**Site testing**

- POC Ab Point-of-care antibody test
- Ag Ab Instrumented antigen/antibody test
- VL Viral load test
- Ab Confirmatory/discriminatory antibody test
- DNA Ultrasensitive DNA test (centralized at JHU)

**HPTN LC testing**

- Ag Ab ARCHITECT antigen/antibody test
- qual RNA APTIMA qualitative RNA test
- Ab Geenius



\*Selected cases



# Extended HPTN LC Testing


## HIV testing

### Back-testing




CAB arm: All visits

TDF/FTC arm: Enrollment, weeks 2, 4, 5



CAB arm: Enrollment plus three visits prior to the first RNA pos visit

TDF/FTC arm: Enrollment plus one visit prior to the first RNA pos visit



If Ag/Ab test reactive



If qualitative RNA test reactive



Selected cases/visits

## HIV genotyping (VL >500 c/mL)

### CAB arm

- All study visits

### TDF/FTC arm

- First HIV positive visit
- First site positive visit

	ARCHITECT antigen/antibody test
	APTIMA qualitative RNA test
	Geenius discriminatory antibody test
	Viral load test
	Single copy RNA test

## Pharmacology testing

### CAB concentrations

#### CAB arm

Plasma [CAB]: all study visits

Plasma [TFV]: baseline infections, step 3 infections

DBS [TFV-DP]: step 3 infections

### TDF/FTC concentrations


Plasma [TFV]: first site pos, first HIV pos, 3 prior visits

DBS [TFV-DP]: first site pos, 1 prior visit





## 13 Incident, 2 baseline Infections: Cabotegravir




- Step 1: Oral CAB lead-in
- Step 2: CAB LA 600 mg IM
- Step 2: CAB LA injection > 2 week overdue
- Step 3: Open-label TDF/FTC
- Step 3: Overdue TDF/FTC dispensation
- Annual follow-up
- %: Percent adherence to oral lead-in
- : CAB LA 600 mg IM
- : Open-label TDF/FTC dispensed
- : HIV-infection
- \*: First site positive






## **13 Incident, 2 baseline Infections: Cabotegravir**






## 13 Incident, 2 baseline Infections: Cabotegravir






## 13 Incident, 2 baseline Infections: Cabotegravir






## 13 Incident, 2 baseline Infections: Cabotegravir






## 13 Incident, 2 baseline Infections: Cabotegravir






## 13 Incident, 2 baseline Infections: Cabotegravir






## 12 Incident, 4 baseline Infections: Cabotegravir






# HIV Incidence: CAB vs. TDF/FTC




CI, confidence interval




**CAB arm, Group A**  
HIV positive at study enrollment



<b>NRTI:</b>	K65R, M184V
<b>NNRTI:</b>	L100I, K103N, P225H
Ag/Ab test	-
Qualitative RNA test	+
Confirmatory Ab test	+
Viral load	4,010




<b>WT</b>	<b>WT</b>	<b>WT</b>	
Ag/Ab test	-	-	
Qualitative RNA test	+	+	
Confirmatory Ab test	+	+	
Viral load	50,080	20,760	700





● CAB concentration    — CAB injection    — First HIV positive visit    — First site positive visit    — First HIV positive visit and first site positive visit  
← # → Weeks between first HIV positive visit and the first site positive visit



The x-axis represents weeks since enrollment. The shaded area represents time on ART.

<b>WT</b>	<b>WT</b>	<b>WT</b>	
Ag/Ab test	-	+	
Qualitative RNA test	+	-	
Confirmatory Ab test	-	-	
Viral load	1,360	<40	ND




<b>WT</b>
Ag/Ab test
Qualitative RNA test
Confirmatory Ab test
Viral load







● CAB concentration    ● CAB injection    — First HIV positive visit    — First site positive visit  
 ← # → Weeks between first HIV positive visit and the first site positive visit



The x-axis represents weeks since enrollment. The shaded area represents time on ART.






● CAB concentration    ━━ CAB injection    — First HIV positive visit    ━━━ First site positive visit    ━━ First HIV positive visit and first site positive visit  
 ← # → Weeks between first HIV positive visit and the first site positive visit


The x-axis represents weeks since enrollment. The shaded area represents time on ART.



**CAB arm, Group B**  
No recent CAB exposure









**CAB arm, Group C**  
Infected during the CAB oral lead-in period




	-	-	-	-	-	-	-	-	-
Ag/Ab test	-	-	-	-	-	-	-	-	-
Qualitative RNA test	-	-	+	+	+	-	+	-	-
Confirmatory Ab test	ND	120	161	137	2,174	1,373	2,549		
Viral load									



	-	-	-	-	-	-	-	-	-
Ag/Ab test	-	-	-	-	-	-	-	-	-
Qualitative RNA test	-	-	+++	++	-	-	-	-	-
Confirmatory Ab test	ND	ND	102,329	<40	375	<40	ND	ND	ND
Viral load									





	-	-	-	-	-	-	-	-	-
Ag/Ab test	-	-	-	-	-	-	-	-	-
Qualitative RNA test	-	-	-	-	-	-	-	-	-
Confirmatory Ab test	-	-	-	-	-	-	-	-	-
Viral load	494	229,810	207,810						



● CAB concentration    — CAB injection    — First HIV positive visit  
— First site positive visit    — First HIV positive visit and first site positive visit  
↔ # Weeks between first HIV positive visit and first site positive visit

The x-axis represents weeks since enrollment. The y-axis represents CAB concentration in mcg/mL. The time on ART.






● CAB concentration    — CAB injection    — First HIV positive visit  
— First site positive visit    — First HIV positive visit and first site positive visit  
↔ # → Weeks between first HIV positive visit and first site positive visit

The x-axis represents weeks since enrollment. The y-axis represents CAB concentration (mcg/mL). The legend indicates CAB concentration (yellow circles), CAB injection (green line), First HIV positive visit (red line), First site positive visit (blue line), and First HIV positive visit and first site positive visit (red dashed line).



**CAB arm, Group D**  
Infected in the setting of on-time CAB  
injections






The shaded area represents time on ART.




Ag/Ab test	-	-	-	-	-	-	-	-	-	+	+	+	+
Qualitative RNA test	-	-	-	-	-	-	+	-	+	-	-	+	+
Confirmatory Ab test								-	-	IND		IND	IND
Viral load					ND	ND					<40		<40

D2



The shaded area represents time on ART.






The shaded area represents time on ART.




## A Cautionary Tale

	Week 0	Week 5	Week 10	Week 15	Week 20	Week 25	Week 30	Week 35	Week 40	Week 45	Week 50	Week 55	Week 60	Week 65
Ag/Ab test	-	-	-	-	-	++			+		+			+
Qualitative RNA test	-	-	-	-	-	++			+		+			+
Confirmatory Ab test					IND	IND			+		+			+
Viral load			<40		158 263		152,730		<40		<40			

D4



The shaded area represents time on ART.






## Key virology findings - CAB arm

- Extended testing identified earlier infection dates in many cases
- Virus loads were often low at the first HIV positive visit
- There was often a prolonged period of viral suppression after infection
- Antibody expression was diminished / delayed in many cases
- In some cases, RNA and Ab tests reverted to negative/non-reactive early in infection






**TDF/FTC arm**





## 39 Incident, 3 Baseline Infections: TDF/FTC





# Key Observations & Conclusions

## Key observations:

- 4 incident infections in the CAB arm occurred despite target plasma CAB concentrations; evaluation of correlates of protection is ongoing
- CAB-LA can delay detection of infection using standard HIV testing algorithms
- INI resistance seen when viremic “escape” occurs at higher CAB concentrations
- INI resistance was not seen in 3 tail-phase infections or 1 tail “escape” case
- 37/39 in the TDF/FTC arm with incident infection had suboptimal or non-adherence

## Conclusions:

- Oral lead-in will be optional in 083 OLE
- Use of VL testing as a primary screen for HIV infection will be assessed in 083 OLE
- In the setting of CAB-LA, prompt diagnosis and ART initiation are needed to prevent resistance

In HPTN 083, CAB-LA and TDF/FTC were both highly effective for prevention of HIV infection. CAB-LA was superior to daily oral TDF/FTC for HIV PrEP in terms of adherence.



# Acknowledgements

## Sponsor

- U.S. National Institute of Allergy and Infectious Diseases (NIAID), National Institute of Mental Health (NIMH), National Institute on Drug Abuse (NIDA), and the National Institute of Child Health and Human Development (NICHD) all components of the U.S. National Institutes of Health (NIH)
- Additional funding from ViiV Healthcare

## HIV Prevention Trials Network (HPTN)

- Laboratory Center (Johns Hopkins University)
- Statistical Center for HIV/AIDS Research and Prevention (SCHARP), Fred Hutchinson Cancer Research Center
- Leadership and Operations Center, FHI 360
- HPTN Leadership

## Pharmaceutical Support

- ViiV Healthcare
- Gilead Sciences, Inc.

Questions? Email [rlandovitz@mednet.ucla.edu](mailto:rlandovitz@mednet.ucla.edu) or



@doc\_in\_a\_box

## HPTN 083 Study Team

**Community Program Managers  
Community Educators & Recruiters,  
CAB Members**

**Our 43 Sites in 7 countries**

**...and our Study Participants!**

